Insights into Internal Structure of a Megabed from Long Distance (130×30 km) Correlation

Fabrizio Felletti , Mattia Marini, Nicolò Bellin contact: fabrizio.felletti@unimi.it

laminae and small-scale hummocky-type different structures and palaeocurrents.

structures. These structures can have quite

Insights into Internal Structure of a Megabed from Long Distance (130×30 km) Correlation: the Contessa Megaturbidite of the Miocene MarnosoArenacea Fm (Serravallian; Northern Apennines, Italy)


Insights into Internal Structure of a Megabed from Long Distance (130×30 km) Correlation


<u>Fabrizio Felletti</u> , Mattia Marini, Nicolò Bellin


INTERNAL DIVISION

Interval 2- Parallel lamination













DISTRIBUTION OF INTERNAL DIVISIONS












Interpretation: These structures can separate different laminasets with be interpreted as combined flow structures. They are evidence of the

Desio' of the University of Milan.



Università degli Studi di Milano, Dipartimento di Scienze della Terra Via Luigi Mangiagalli, 34 - 20133 Milano MI, Italy

depocenters separated by the Verghereto High. To the SE, the thickness of the MA decreases

Via Luigi Mangiagalli, 34 - 20133 Milano MI, Italy

Jniversità degli Studi di Milano, Dipartimento di Scienze della Terra ia Luigi Mangiagalli, 34 - 20133 Milano MI, Italy